

# Asymptotic analysis of powers of matrices

Diego Dominici \*

Department of Mathematics  
State University of New York at New Paltz  
75 S. Manheim Blvd. Suite 9  
New Paltz, NY 12561-2443  
USA

Phone: (845) 257-2607

Fax: (845) 257-3571

## Abstract

We analyze the representation of  $A^n$  as a linear combination of  $A^j$ ,  $0 \leq j \leq k-1$ , where  $A$  is a  $k \times k$  matrix. We obtain a first order asymptotic approximation of  $A^n$  as  $n \rightarrow \infty$ , without imposing any special conditions on  $A$ .

We give some examples showing the application of our results.

Keywords: Matrix powers, asymptotic approximations of integrals, generating functions.

MSC-class: 15A15 (Primary) 34E05 (Secondary)

## 1 Introduction

In a recent article [1], Abu-Saris and Ahmad showed how to compute the powers of a matrix without having to compute its eigenvalues. Their main result was:

---

\*e-mail: dominicd@newpaltz.edu

**Theorem 1** *If  $A$  is a  $k \times k$  matrix with characteristic polynomial*

$$P(x) = x^k + \sum_{j=0}^{k-1} a_j x^j, \quad (1)$$

*then,*

$$A^n = \sum_{j=0}^{k-1} b_j(n) A^j, \quad n \geq k \quad (2)$$

*where*

$$\begin{aligned} b_j(k) &= -a_j, \quad 0 \leq j \leq k-1, \quad b_{-1}(n) = 0, \quad n \geq k, \\ b_j(n+1) &= b_{j-1}(n) - a_j b_{k-1}(n), \quad n \geq k, \quad 0 \leq j \leq k-1. \end{aligned} \quad (3)$$

The purpose of this paper is to find an asymptotic representation for the numbers  $b_j(n)$  as  $n \rightarrow \infty$ , which using (2) will give an asymptotic representation of  $A^n$  for large  $n$ . Since the coefficients  $b_j(n)$  depend only on  $P(x)$ , our estimates will be valid for similar matrices.

The asymptotic behavior of powers of matrices has been considered before by other authors. In [5] and [6], Gautschi computed upper bounds for  $A^n$  and  $\|A^n\|$ , where  $\|A\|$  is a norm of  $A$ . Estimates of  $\|A^n\|$  were also studied in [2], [3], [11], [13] and [14].

In [4], Friedland and Schneider considered the matrix

$$B^{(m)} = A^m (I + \dots + A^{q-1}), \quad m \geq 1$$

where  $A$  is a nonnegative matrix and  $q$  is a certain positive integer. They proved a theorem on the growth of  $B^{(m)}$  under the assumption that the spectral radius of  $A$  is equal to one. Powers of nonnegative matrices were also analyzed by Lindqvist in [8]. Rothblum [10], obtained Cesaro asymptotic expansions of  $\sum_{i=0}^N A^i$ , where  $A$  is a complex matrix with spectral radius less than or equal to one.

This paper is organized as follows: In Section 2 we find an integral representation for the exponential generating function  $G_j(z)$  of the coefficients  $b_j(n)$ . We obtain exact formulas for  $G_j(z)$  and  $b_j(n)$  in the special case of the matrix  $A$  having  $k$  distinct eigenvalues. We conclude the section with some examples.

In Section 3 we give an exact representation and a first order asymptotic approximation for  $b_j(n)$ , as  $n \rightarrow \infty$ . We consider the cases of simple and multiple eigenvalues. Our formulas are relatively easy to implement and offer very accurate estimates of  $b_j(n)$ , and therefore of  $A^n$ , for large  $n$ . We present some examples for different cases of  $P(x)$ .

## 2 Generating function

In this section we shall find an exponential generating function for the coefficients  $b_j(n)$ . First, let us define the spectral radius  $\rho(A)$  of the matrix  $A$  by

$$\rho(A) = \max \{ |\lambda| \mid P(\lambda) = 0 \}. \quad (4)$$

**Theorem 2** *Let  $G_j(z)$  be defined by*

$$G_j(z) = \sum_{n \geq 0} b_j(n+k) \frac{z^n}{n!}, \quad (5)$$

*Then, we have*

$$G_j(z) = -\frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} s^{k-j-1} \frac{p_j(s)}{P(s)} e^{zs} ds, \quad (6)$$

*where  $c > \rho(A)$ ,*

$$p_j(s) = \sum_{l=0}^j a_l s^l, \quad 0 \leq j \leq k-1, \quad (7)$$

*and  $P(s)$  is the characteristic polynomial of  $A$  defined in (7).*

**Proof.** If we use (5) in (3), we obtain

$$G'_j = G_{j-1} - a_j G_{k-1}, \quad G_j(0) = -a_j, \quad 0 \leq j \leq k-1 \quad (8)$$

with  $G_{-1}(z) = 0$ . Taking the Laplace transform of  $G_j(z)$ ,

$$L_j(s) = \int_0^\infty G_j(z) e^{-zs} dz$$

in (8) we get

$$sL_j + a_j = L_{j-1} - a_j L_{k-1}, \quad 0 \leq j \leq k-1 \quad (9)$$

and  $L_{-1}(z) = 0$ . ■

The solution of (9) is given by

$$L_j(s) = -s^{k-j-1} \frac{p_j(s)}{P(s)}, \quad 0 \leq j \leq k-1 \quad (10)$$

where

$$p_j(s) = \sum_{l=0}^j a_l s^l, \quad 0 \leq j \leq k-1,$$

and  $p_{-1}(s) = 0$ . Inverting the Laplace transform in (10) the theorem follows.

**Remark 3** *Since*

$$\lim_{|s| \rightarrow \infty} L_j(s) = 0, \quad 0 \leq j \leq k-1$$

*we can replace the Bromwich contour in (6) with a circle  $\mathcal{C}$  of radius  $R$  centered at the origin [7]*

$$G_j(z) = -\frac{1}{2\pi i} \int_{\mathcal{C}} s^{k-j-1} \frac{p_j(s)}{P(s)} e^{zs} ds \quad (11)$$

*with  $R > \rho(A)$ .*

**Corollary 4** *If*

$$P(s) = (s - \lambda_1)(s - \lambda_2) \cdots (s - \lambda_k),$$

*where the eigenvalues  $\lambda_i$  are all distinct, then*

$$G_j(z) = -\sum_{l=1}^k (\lambda_l)^{k-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)} \exp(\lambda_l z) \quad (12)$$

*and*

$$b_j(n) = -\sum_{l=1}^k (\lambda_l)^{n-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)}. \quad (13)$$

**Proof.** Applying the residue theorem to (11) we obtain

$$G_j(z) = - \sum_{P(\lambda)=0} \operatorname{Res} \left[ s^{k-j-1} \frac{p_j(s)}{P(s)} e^{zs}; \lambda \right] \quad (14)$$

which in turn gives (12) after computing

$$\begin{aligned} \operatorname{Res} \left[ s^{k-j-1} \frac{p_j(s)}{P(s)} e^{zs}; \lambda_l \right] &= \lim_{s \rightarrow \lambda_l} s^{k-j-1} p_j(s) e^{zs} \frac{(s - \lambda_l)}{P(s)} \\ &= \lim_{s \rightarrow \lambda_l} s^{k-j-1} p_j(s) e^{zs} \frac{1}{P'(s)} = (\lambda_l)^{k-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)} \exp(\lambda_l z). \end{aligned}$$

Writing (12) as

$$G_j(z) = - \sum_{l=1}^k (\lambda_l)^{k-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)} \sum_{n \geq 0} (\lambda_l)^n \frac{z^n}{n!}$$

and changing the order of summation, we have

$$G_j(z) = \sum_{n \geq 0} \left[ - \sum_{l=1}^k (\lambda_l)^{n+k-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)} \right] \frac{z^n}{n!}$$

which implies

$$b_j(n+k) = - \sum_{l=1}^k (\lambda_l)^{n+k-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)}.$$

■

**Example 5** In [1] the authors considered the following examples:

1.

$$P(x) = x^3 - 7x^2 + 16x - 12 = (x-2)^2(x-3).$$

Using (10) we have

$$\begin{aligned} L_0(s) &= \frac{-s^2(12)}{(s-2)^2(s-3)} \\ L_1(s) &= \frac{-s(16s-12)}{(s-2)^2(s-3)} \\ L_2(s) &= \frac{-(-7s+16s-12)}{(s-2)^2(s-3)} \end{aligned}$$

and inverting we obtain

$$\begin{aligned} G_0(z) &= -12(8 + 4z)e^{2z} + 108e^{3z} \\ G_1(z) &= 4(23 + 10z)e^{2z} - 108e^{3z} \\ G_2(z) &= -4(5 + 2z)e^{2z} + 27e^{3z}. \end{aligned}$$

Expanding in series we get

$$G_0(z) = -96 \sum_{n \geq 0} 2^n \frac{z^n}{n!} - 48 \sum_{n \geq 0} 2^{n-1} n \frac{z^n}{n!} + 108 \sum_{n \geq 0} 3^n \frac{z^n}{n!}$$

and from (5) we conclude that

$$b_0(n+3) = -96 \times 2^n - 48 \times 2^{n-1}n + 108 \times 3^n$$

or

$$\begin{aligned} b_0(n) &= -96 \times 2^{n-3} - 48 \times 2^{n-4}(n-3) + 108 \times 3^{n-3} \\ &= -3(1+n) \times 2^n + 4 \times 3^n. \end{aligned}$$

Similar calculations give

$$\begin{aligned} b_1(n) &= \left(4 + \frac{5}{2}n\right) \times 2^n - 4 \times 3^n \\ b_2(n) &= -\left(1 + \frac{1}{2}n\right) \times 2^n + 3^n, \end{aligned}$$

in agreement with the results shown in [1].

2.

$$P(x) = x^3 - 5x^2 + 6x = x(x-2)(x-3).$$

We can apply (13) directly and obtain

$$\begin{aligned} b_0(n) &= 0 \\ b_1(n) &= \frac{3}{2} \times 2^n - \frac{2}{3} \times 3^n \\ b_2(n) &= -\frac{1}{2} \times 2^n + \frac{1}{3} \times 3^n. \end{aligned}$$

3.

$$P(x) = x^5 - 5x^4 + 10x^3 - 20x^2 - 15x - 4 = (x - 4)(x^4 - x^3 + 6x^2 + 4x + 1).$$

Although (as the authors noted) MAPLE is unable to compute the zeros of  $P(x)$  exactly, it can provide us with very accurate numerical approximations

$$\begin{aligned}\lambda_1 &= 4 \\ \lambda_2 &= 0.8090169944 + 2.489898285i \\ \lambda_3 &= 0.8090169944 - 2.489898285i \\ \lambda_4 &= -0.3090169944 + 0.2245139883i \\ \lambda_5 &= -0.3090169944 - 0.2245139883i\end{aligned}$$

which we can use in (13) to get

$$b_j(n) = \frac{C_j}{305} 4^n - \sum_{l=2}^5 (\lambda_l)^{n-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)}$$

with

$$C_0 = 1, C_1 = 4, C_2 = 6, C_3 = -1, C_4 = 1.$$

Note that, for  $0 \leq j \leq 4$ , we have

$$\sum_{l=2}^5 (\lambda_l)^{n-j-1} \frac{p_j(\lambda_l)}{P'(\lambda_l)} = O(|\lambda_2|^n) = O(2.618^n)$$

as  $n \rightarrow \infty$ .

**Example 6** Let  $A$  be the matrix

$$A = \begin{pmatrix} 1 & 2 \\ -1 & -1 \end{pmatrix}$$

with characteristic polynomial

$$P(x) = x^2 + 1 = (x - i)(x + i).$$

Using (13) we have

$$\begin{aligned}b_0(n) &= \cos\left(\frac{\pi}{2}n\right) \\ b_1(n) &= \sin\left(\frac{\pi}{2}n\right)\end{aligned}$$

and from (2) we get

$$A^n = \begin{pmatrix} \cos\left(\frac{\pi}{2}n\right) + \sin\left(\frac{\pi}{2}n\right) & 2\sin\left(\frac{\pi}{2}n\right) \\ -\sin\left(\frac{\pi}{2}n\right) & \cos\left(\frac{\pi}{2}n\right) - \sin\left(\frac{\pi}{2}n\right) \end{pmatrix}.$$

In particular, we have

$$A^n = \begin{cases} I, & n \equiv 0(4) \\ A, & n \equiv 1(4) \\ -I, & n \equiv 2(4) \\ -A, & n \equiv 3(4) \end{cases}$$

where  $I$  denotes the identity matrix.

**Example 7** This example appeared in [9]. Let  $A$  be the matrix

$$A = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}$$

with characteristic polynomial

$$P(x) = x^2 - x - 1 = (x - \alpha)(x - \beta),$$

where

$$\alpha = \frac{1}{2}(1 + \sqrt{5}), \quad \beta = \frac{1}{2}(1 - \sqrt{5}).$$

Then, from (13), we have

$$\begin{aligned} b_0(n) &= \frac{1}{\sqrt{5}}(\alpha^{n-1} - \beta^{n-1}) = f_{n-1} \\ b_1(n) &= \frac{1}{\sqrt{5}}(\alpha^n - \beta^n) = f_n \end{aligned}$$

where  $f_n$  is the  $n$ th Fibonacci number. Thus,

$$A^n = \begin{pmatrix} f_n + f_{n-1} & f_n \\ f_n & f_{n-1} \end{pmatrix} = \begin{pmatrix} f_{n+1} & f_n \\ f_n & f_{n-1} \end{pmatrix}.$$



### 3 Asymptotic analysis

We begin by finding an integral representation of the coefficients  $b_j(n)$ .

**Lemma 8** *The numbers  $b_j(n)$  can be represented as*

$$b_j(n) = -\frac{1}{2\pi i} \int_{\mathcal{C}} s^{n-j-1} \frac{p_j(s)}{P(s)} ds \quad (15)$$

where  $\mathcal{C}$  is a circle of radius  $R > \rho(A)$  centered at the origin and the polynomials  $p_j(s)$  were defined in (7).

**Proof.** Since the power series

$$e^{zs} = \sum_{n \geq 0} s^n \frac{z^n}{n!}$$

converges uniformly on  $|s| \leq R$ , we can interchange integration and summation in (11) and obtain

$$G_j(z) = \sum_{n \geq 0} \left[ -\frac{1}{2\pi i} \int_{\mathcal{C}} s^{n+k-j-1} \frac{p_j(s)}{P(s)} ds \right] \frac{z^n}{n!}.$$

Then, (5) implies

$$b_j(k+n) = -\frac{1}{2\pi i} \int_{\mathcal{C}} s^{n+k-j-1} \frac{p_j(s)}{P(s)} ds, \quad 0 \leq j \leq k-1$$

and the result follows. ■

**Remark 9** *An alternative method for approximating the coefficients  $b_j(n)$  is to write (15) as*

$$b_j(n) = -\frac{R^{n-j}}{2\pi} \int_0^{2\pi} \exp[it(n-j)] \frac{p_j(Re^{it})}{P(Re^{it})} dt \quad (16)$$

with  $R > \rho(A)$  and to compute the integral (16) numerically. This approach offers the advantage of avoiding the computation of the eigenvalues of  $A$ .

We now have all the necessary elements to establish our main theorem.

**Theorem 10** *Let*

$$\rho(A) = |\lambda| > |\lambda_2| > \cdots > |\lambda_r|$$

*be the eigenvalues of the matrix  $A$ , i.e.,*

$$P(x) = (x - \lambda)^m (x - \lambda_2)^{m_2} \cdots (x - \lambda_r)^{m_r} \quad (17)$$

*with  $r \leq k$ . Then,*

$$b_j(n) \sim -\lambda^{n-m-j} \frac{p_j(\lambda)}{P^{(m)}(\lambda)} m! \binom{n-k}{m-1}, \quad n \rightarrow \infty, \quad (18)$$

*where*

$$P^{(m)}(\lambda) = \left. \frac{d^m P}{ds^m} \right|_{s=\lambda}.$$

**Proof.** To find an asymptotic approximation of (15), we shall use a modified version of Darboux's Method [12]. We write

$$s^{n-j-1} \frac{p_j(s)}{P(s)} = s^{n-k} \times \frac{s^{k-j-1} p_j(s)}{P(s)},$$

so that  $\deg(s^{k-j-1} p_j) = k-1$  and  $\deg(P) = k$ .

From (17) we have

$$\frac{s^{k-j-1} p_j(s)}{P(s)} \sim \frac{\lambda^{k-j-1} p_j(\lambda)}{(s-\lambda)^m g(\lambda)}, \quad s \rightarrow \lambda \quad (19)$$

where

$$g(x) = (x - \lambda_2)^{m_2} \cdots (x - \lambda_r)^{m_r}.$$

Using the Binomial Theorem, we obtain

$$s^{n-k} = \sum_{l=0}^{n-k} (s-\lambda)^l \binom{n-k}{l} \lambda^{n-k-l}. \quad (20)$$

Combining (19) and (20), we get

$$s^{n-j-1} \frac{p_j(s)}{P(s)} \sim \lambda^{n-m-j} \frac{p_j(\lambda)}{g(\lambda)} \binom{n-k}{m-1} \frac{1}{(s-\lambda)}, \quad s \rightarrow \lambda$$

and therefore

$$b_j(n) = -\frac{1}{2\pi i} \int_c s^{n-j-1} \frac{p_j(s)}{P(s)} ds \sim -\lambda^{n-m-j} \frac{p_j(\lambda)}{g(\lambda)} \binom{n-k}{m-1}, \quad n \rightarrow \infty. \quad (21)$$

To find the value of  $g(\lambda)$ , we use L'Hopital's Theorem

$$g(\lambda) = \lim_{s \rightarrow \lambda} \frac{P(s)}{(s-\lambda)^m} = \lim_{s \rightarrow \lambda} \frac{P^{(m)}(s)}{m!} = \frac{P^{(m)}(\lambda)}{m!}. \quad (22)$$

Replacing (22) in (21), we obtain (18). ■

**Remark 11** *Note that when  $m = 1$  we recover the leading term in (13).*

If more than one eigenvalue has absolute value equal to the spectral radius of  $A$ , the asymptotic behavior of  $b_j(n)$  can be obtained by adding the contributions from each eigenvalue. We state this formally in the following corollary.

**Corollary 12** *If*

$$\rho(A) = |\lambda_1| = |\lambda_2| = \dots = |\lambda_r|,$$

*with respective multiplicities  $m_1, m_2, \dots, m_r$ , then*

$$b_j(n) \sim -\sum_{l=1}^r (\lambda_l)^{n-m_l-j} \frac{p_j(\lambda_l)}{P^{(m_l)}(\lambda_l)} (m_l)! \binom{n-k}{m_l-1}, \quad n \rightarrow \infty. \quad (23)$$

**Remark 13** *Since*

$$\binom{n-k}{m_l-1} \sim \frac{1}{(m_l-1)!} n^{m_l-1}, \quad n \rightarrow \infty$$

*we have*

$$b_j(n) \sim -\sum_{l=1}^r (\lambda_l)^{n-m_l-j} \frac{p_j(\lambda_l)}{P^{(m_l)}(\lambda_l)} m_l n^{m_l-1}, \quad n \rightarrow \infty.$$

*Therefore, in the case of several eigenvalues located on the circle  $|s| = \rho(A)$ , the dominant term in (23) will correspond to the eigenvalue with the greatest multiplicity.*

**Example 14** In Example 5 (1) we consider

$$P(x) = x^3 - 7x^2 + 16x - 12 = (x - 2)^2 (x - 3).$$

In this case,  $\lambda = 2$ ,  $m = 2$  and  $k = 3$ . From (18), we get

$$\begin{aligned} b_0(n) &\sim 4 \times 3^n \\ b_1(n) &\sim -4 \times 3^n \\ b_2(n) &\sim 3^n \end{aligned}$$

which are the leading terms in the solution previously obtained.

**Example 15** We now consider the case of more than one eigenvalue having absolute value equal to  $\rho(A)$ . Let

$$P(x) = x^4 + x^3 - 15x^2 - 9x + 54 = (x - 2)(x - 3)(x + 3)^2.$$

In this case,  $\lambda_1 = -3$ ,  $m_1 = 2$ ,  $\lambda_2 = 3$ ,  $m_2 = 1$  and  $k = 4$ . From (23), we have

$$\begin{aligned} b_0(n) &\sim -\frac{1}{5}n(-3)^n + \frac{4}{5}(-3)^n - \frac{1}{2}3^n \\ b_1(n) &\sim \frac{1}{10}n(-3)^n - \frac{2}{5}(-3)^n - \frac{1}{12}3^n \\ b_2(n) &\sim \frac{1}{45}n(-3)^n - \frac{4}{45}(-3)^n + \frac{1}{9}3^n \\ b_3(n) &\sim -\frac{1}{90}n(-3)^n + \frac{2}{45}(-3)^n + \frac{1}{36}3^n. \end{aligned}$$

The exact values are

$$\begin{aligned} b_0(n) &= -\frac{1}{5}n(-3)^n + \frac{21}{50}(-3)^n - \frac{1}{2}3^n + \frac{27}{25}2^n \\ b_1(n) &= \frac{1}{10}n(-3)^n - \frac{83}{300}(-3)^n - \frac{1}{12}3^n + \frac{9}{25}2^n \\ b_2(n) &= \frac{1}{45}n(-3)^n + \frac{2}{225}(-3)^n + \frac{1}{9}3^n - \frac{3}{25}2^n \\ b_3(n) &= -\frac{1}{90}n(-3)^n + \frac{11}{900}(-3)^n + \frac{1}{36}3^n - \frac{1}{25}2^n. \end{aligned}$$

As we observed before, the main contribution comes from the eigenvalue of maximum multiplicity, in this case  $\lambda_1 = -3$ .

**Example 16** Finally, let's consider the case of complex eigenvalues of multiplicity greater than one located on the circle  $|s| = \rho(A)$ . Let

$$\begin{aligned} P(x) &= x^5 - 9x^4 + 34x^3 - 66x^2 + 65x - 25 \\ &= (x - 1) [x - (2 + i)]^2 [x - (2 - i)]^2. \end{aligned}$$

In this case,  $\lambda_1 = 2 + i$ ,  $m_1 = 2$ ,  $\lambda_2 = 2 - i$ ,  $m_2 = 2$  and  $k = 5$ . From (23), we obtain

$$\begin{aligned} b_0(n) &\sim \frac{1}{4} \left( \sqrt{5} \right)^n (n - 5) [\cos(\theta n) - 7 \sin(\theta n)] \\ b_1(n) &\sim -\frac{1}{10} \left( \sqrt{5} \right)^n (n - 5) [2 \cos(\theta n) - 39 \sin(\theta n)] \\ b_2(n) &\sim -\frac{1}{10} \left( \sqrt{5} \right)^n (n - 5) [2 \cos(\theta n) + 31 \sin(\theta n)] \\ b_3(n) &\sim \frac{1}{10} \left( \sqrt{5} \right)^n (n - 5) [2 \cos(\theta n) + 11 \sin(\theta n)] \\ b_4(n) &\sim -\frac{1}{20} \left( \sqrt{5} \right)^n (n - 5) [\cos(\theta n) + 3 \sin(\theta n)] \end{aligned}$$

with

$$\theta = \arctan \left( \frac{1}{2} \right).$$

The exact values are

$$\begin{aligned} b_0(n) &= \frac{1}{4} \left( \sqrt{5} \right)^n [(n - 21) \cos(\theta n) + (-7n + 22) \sin(\theta n)] + \frac{25}{4} \\ b_1(n) &= -\frac{1}{10} \left( \sqrt{5} \right)^n [2(n - 50) \cos(\theta n) + (-39n + 125) \sin(\theta n)] - 10 \\ b_2(n) &= -\frac{1}{10} \left( \sqrt{5} \right)^n [(2n + 65) \cos(\theta n) + (31n - 100) \sin(\theta n)] + \frac{13}{2} \\ b_3(n) &= \frac{1}{10} \left( \sqrt{5} \right)^n [2(n + 10) \cos(\theta n) + (11n - 35) \sin(\theta n)] - 2 \\ b_4(n) &= -\frac{1}{20} \left( \sqrt{5} \right)^n [(n + 5) \cos(\theta n) + (3n - 10) \sin(\theta n)] + \frac{1}{4}. \end{aligned}$$

## References

- [1] R. Abu-Saris and W. Ahmad. Avoiding eigenvalues in computing matrix powers. *Amer. Math. Monthly*, 112(5):450–454, 2005.

- [2] N. Borovikh, D. Drissi, and M. N. Spijker. A bound on powers of linear operators, with relevance to numerical stability. *Appl. Math. Lett.*, 15(1):47–53, 2002.
- [3] N. Borovikh and M. N. Spijker. Resolvent conditions and bounds on the powers of matrices, with relevance to numerical stability of initial value problems. *J. Comput. Appl. Math.*, 125(1-2):41–56, 2000.
- [4] S. Friedland and H. Schneider. The growth of powers of a nonnegative matrix. *SIAM J. Algebraic Discrete Methods*, 1(2):185–200, 1980.
- [5] W. Gautschi. The asymptotic behaviour of powers of matrices. *Duke Math. J.*, 20:127–140, 1953.
- [6] W. Gautschi. The asymptotic behaviour of powers of matrices. II. *Duke Math. J.*, 20:375–379, 1953.
- [7] W. R. LePage. *Complex variables and the Laplace transform for engineers*. Dover Publications Inc., New York, 1980.
- [8] B. H. Lindqvist. Asymptotic properties of powers of nonnegative matrices, with applications. *Linear Algebra Appl.*, 114/115:555–588, 1989.
- [9] K. H. Rosen. *Elementary number theory and its applications*. Addison-Wesley, Reading, MA, fourth edition, 2000.
- [10] U. G. Rothblum. Expansions of sums of matrix powers. *SIAM Rev.*, 23(2):143–164, 1981.
- [11] P. Stavinoha. Convergence of  $L_p$ -norms of a matrix. *Apl. Mat.*, 30(5):351–360, 1985.
- [12] R. Wong. *Asymptotic approximations of integrals*, volume 34 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2001.
- [13] N. J. Young. Analytic programmes in matrix algebras. *Proc. London Math. Soc. (3)*, 36(2):226–242, 1978.
- [14] N. J. Young. Norms of powers of matrices with constrained spectra. *Linear Algebra Appl.*, 23:227–244, 1979.